

Beyond Breakthrough: Optimizing the Breakthrough Therapy Designation

Pre-Conference Draft

Introduction:

Advances in our understanding of disease processes, genetics, manufacturing technologies, and innovative clinical trial design have enabled the development of novel therapeutic agents for the treatment of patients with cancer. In oncology, the ability to target a novel agent against a driver oncogene or protective immune checkpoint has led to several therapeutic breakthroughs in diseases with limited or no systemic treatment options. These breakthroughs have established new classes of therapeutics leading to, in some instances, unprecedented improvements in clinical outcomes for patients with cancer.

Regulatory review processes are time and resource intensive for drug sponsors and the United States Food and Drug Administration (FDA). The FDA leverages several tools to safely and efficiently facilitate development and review of agents intended for treatment of patients with life-threatening conditions without compromising the rigorous standards established for their approval.

Breakthrough Therapy Designation (BTD) facilitates the efficient development of both drugs and biologics (hereafter referred to as “drugs”) intended to treat serious or life-threatening illnesses for which there is preliminary clinical evidence demonstrating that the investigational therapy may offer substantial improvement on a clinically significant endpoint(s) over available therapies.¹ BTD pro-

Objectives

Characterize the rate-limiting steps and challenges encountered throughout the development of oncology products with Breakthrough Therapy Designation (BTD).

Propose recommendations that address commonly encountered challenges and identify best practices to ensure development programs maximize the benefits of BTD.

Delineate key topics and optimal timing for interactions with the FDA when requesting BTD (pre-BTD) and after receiving BTD (post-BTD).

Acknowledgements

We thank the following organizations and the many representatives for their commitment of time, expertise, and insights to identify actionable opportunities to optimize the use of Breakthrough Therapy Designation.

Amgen
Astellas Pharma
Bayer
Bluebird Bio
Bristol Myers Squibb
Deciphera
EMD Serono
Eli Lilly
Genentech
Iovance Biotherapeutics
Janssen
Kyowa Kirin
Loxo Oncology at Lilly
Merck & Co., Inc.
MorphoSys
Nektar Therapeutics
Novartis
Pfizer
Seagen
Takeda
U.S. Food and Drug Administration (FDA)

This whitepaper should not be construed to represent FDA's views or policies and does not represent official FDA guidance.

vides sponsors with early opportunities for FDA interaction and enhanced guidance, including proactive organizational commitment and coordination involving senior FDA managers and experienced regulatory project management staff. Additionally, BTM often offers a pathway to eligibility for rolling or priority review.²

Since 2012, the number of BTM requests each year has increased. To date, the FDA has received over one thousand requests for BTM and granted more than four hundred requests.³ Both FDA and commercial sponsors prioritize internal resources to help ensure that the most promising products receive BTM and undergo clinical development as efficiently as possible without compromising safety, efficacy, or quality. As a result of this work, BTM has facilitated timely development and approval of 205 products, 61% of which are oncology products. The program has been particularly successful in getting safe and effective novel treatments approved for patients with cancer, particularly new treatments that have not been previously approved for other indications. It is estimated that BTM has shortened the time from IND submission to approval for 60 original applications (not previously approved) for oncology products by a median of 2.3 years compared to oncology products without BTM.⁴

Given the breadth of experience using BTM in oncology over the past eight years, a multistakeholder working group conducted a landscape analysis to identify opportunities to improve implementation of the BTM program in oncology. Evaluating the successful use of the BTM program and challenges associated with optimal use of BTM in oncology will help inform strategies for sustaining its impact on drug development. Historically, much of the focus on BTM has been on the qualifying criteria and processes leading to receipt of BTM; now is the time to identify pressure points and best practices, for both commercial sponsors and the FDA, to optimize implementation of the program within the evolving landscape of oncology drug development to support efficient, successful development and approval of new safe and effective cancer treatments.

Friends of Cancer Research (*Friends*) conducted a survey to solicit input from over 20 commercial sponsors that varied in company size and the range of experiences in terms of number of therapies that have received BTM and approved therapies with BTM to identify challenges and formulate recommendations for optimizing the use of BTM. All sponsors noted the positive impact of BTM in oncology drug development. Several key areas to further optimize use of BTM emerged.

- 1. Clarify expectations for necessary evidence to receive BTM:** Sponsors may find it challenging to anticipate when and what to submit with preliminary and formal BTM requests. There is also a need for additional clarity about the types and quantity of evidence needed to support BTM, particularly with respect to early preliminary clinical evidence in oncology.
- 2. Enhance communication between sponsors and FDA:** The opportunity for enhanced interactions with FDA provided by BTM support proactive identification and resolution of issues; however, at certain stages of development, such as pre-BTM or between milestone meetings, there may be additional opportunities to streamline clinical development through sponsor-FDA interaction.



- 3. Support inter-disciplinary coordination and improve transparency:** In oncology, many drug development programs are increasingly complex and require close coordination between multiple disciplines, and often between Offices and Centers at the FDA (e.g., applications with BTM involving complimentary or companion diagnostics, employing a novel platform or endpoint, targeting rare diseases, and/or incorporating innovative trial designs). Additional guidance on how best to strengthen coordination between Centers, when relevant, at the time of BTM may be beneficial to ensure complexity does not lead to delays.
- 4. Provide additional support to address rate-limiting steps in drug development:** BTM facilitates both clinical and CMC aspects of development; however, challenges can occur during drug development, particularly for novel or innovative technology platforms. CMC development issues can be the rate-limiting step in drug development and approval, and flexibility and timely interactions between FDA and sponsors can be crucial to identify and resolve these issues to mitigate delays. Dose selection and justification is also an important component of drug development and can be a challenge with an expedited drug development timeline.

After conducting the survey, *Friends* convened multiple focus groups with key stakeholders, including the FDA, to identify potential practical solutions in these key areas to support optimal use of BTM.

Opportunity 1. Optimize the timing of BTM and improve communication on expectations for data necessary to receive and maintain BTM.

Provide additional clarity regarding the criteria for BTM to optimize the timing for submission of a BTM request.

Because BTM confers the opportunity for enhanced interactions between sponsors and a multi-disciplinary FDA team including senior FDA staff, and since critical aspects of drug development such as dose optimization and manufacturing can be rate-limiting, BTM has the most potential to positively impact development of drugs that have not previously received FDA approval for another indication. Similarly, the timing of BTM is critical; for example, when BTM is granted based on top-level results of a pivotal trial that will provide the primary evidence to support a marketing application, there may be limited potential for enhanced interactions to result in meaningful improvements to drug development, as opposed to when BTM is granted prior to initiation of a registrational trial.

Among sponsors, there can be uncertainty regarding the level of clinical evidence required to meet the qualifying criteria for BTM. Inclusion of annotation in the preliminary BTM teleconference template to describe the type and scope of preliminary evidence that are generally needed to support a BTM request submission for an oncology product could help facilitate preparation of documents and meaningful preliminary BTM teleconference discussions, while also reducing the number of preliminary BTM teleconferences that clearly lack sufficient data. Additionally, it may be beneficial if sponsors apply for BTM at a time when there is sufficient data to meet the qualifying criteria for BTM but early enough to fully leverage the enhanced interactions provided by

BTD, ideally no later than the time of completion of Phase 2 development. Sponsors may benefit from routine use of preliminary BTD advice teleconferences to discuss eligibility of requests for BTD and gain a better understanding regarding the appropriate timeline and data package necessary to support a BTD request. An oncology-specific guidance describing general guidelines for preparing for a preliminary BTD discussion, content of a BTD request submission, and efficacy considerations for meeting the criteria for BTD may be beneficial.

Clarify procedures and decision-making regarding withdrawing or rescinding BTD and better understand its downstream impact on development/approval to support integrity of the program.

If a program no longer qualifies for BTD, the sponsor can voluntarily withdraw their BTD, or FDA can rescind it. It is sometimes unclear what the timepoints are for re-evaluating the status of BTD or common reasons for withdrawal. Transparent communication regarding the considerations and procedures used by FDA to evaluate whether BTD should be rescinded or withdrawn could ensure a designation is robust and fair. There are context-dependent considerations, such as the timing of the receipt of BTD, stage of development of the program when BTD is withdrawn, and the status of other available therapies, which may impact the public messaging around why withdrawal occurred. Prior to the voluntary withdrawal or rescinding of BTD, the sponsor and the FDA could engage in discussions surrounding the rationale for a planned withdrawal or rescinding of BTD and the implications of these actions on the drug development program. Sponsors may also benefit from an explanation of how withdrawal impacts a program's ability to participate in other Oncology Center for Excellence (OCE) pilots (e.g., Real Time Oncology Review, Project Orbis).

Opportunity 2. Improve mechanisms to enable meaningful discussions between FDA and sponsors that clearly align with key decision points.

Ensure productive and timely discussions between the FDA and sponsor.

It is beneficial to have post-BTD discussions as early as possible so the drug program can maximize BTD's enhanced opportunities for interaction and advice. Ideally, sponsors should prepare for and request the comprehensive post-BTD multidisciplinary meeting so that it occurs in a timely fashion (within the first six months of receiving BTD).⁵ During the meeting, it may be helpful to discuss and agree on a high-level communication plan and estimated timeline for future interactions that address specific drug development aspects based upon anticipated milestones while maintaining flexibility.

Meetings following the comprehensive post-BTD multidisciplinary meeting could be focused to align on the specific needs of a drug development program at each stage of development. The benefit of collaborative discussions with the FDA and sponsors may be more fully realized when questions are focused on a single discrete issue. This could also shorten the lead time for the meeting and reduce the burden on both the FDA and the sponsor to prepare for the meeting. If questions arise between formal meetings, meeting requests targeted to address specific topics could be considered. PDUFA VII goals include a proposed Type D meeting that would be suit-



able to address a narrow set of issues.⁶ Specifically, Type D meetings are intended to address a follow up issue after a formal meeting, a narrow issue the sponsor would like FDA input on, or a general question that does not require detailed advice. During focus group discussions, participants suggested topics such as CMC/product quality-related hurdles, trial design-related issues, and timing of dose optimization studies for meetings like Type D meetings. Timelines for drug development may need to be coordinated when the program is expedited, and key questions that may be rate-limiting may not necessarily arise in alignment with the timing of traditional milestone meetings for aspects such as CMC and clinical considerations. The development of a mechanism to update FDA on key components of drug development could help identify when meetings might be valuable to head off potential rate-limiting obstacles to oncology drug development.

Provide additional clarity to ensure a better understanding of which types of meetings are optimal for specific aspects of drug development for products with BTD.

Formal meetings such as Type B meetings are generally held within 60 days of their request (70 days for end-of-phase meetings) and require extensive preparation on the part of sponsors and FDA staff. As such, these meetings may not always be amenable to post-BTD drug development timeframes, therefore additional meeting strategies are likely to be needed for BTD product development. Table 1 outlines available and proposed meeting types to support development of drugs with BTD. Strategies to formally integrate and operationalize issue specific meetings for products with BTD could be informed through pilot projects such as the Complex Innovative Trial Design Pilot Meeting Program (CID) and Model-Informed Drug Development (MIDD) Pilot Program. Given the fast pace of development post-BTD, decisions are made decisively and quickly, and timely interactions are extremely important.

In addition to timely discussions, it may help to have enhanced interactions and feedback from the FDA. Proactive and thoughtful interactions support expedited development and review of products with BTD. Optimizing meeting structure and approach could enable discussions between FDA and sponsors to occur more frequently, help address issues earlier in development, and provide opportunities for proactive planning of manufacturing and testing strategies and clinical development relative to traditional drug development approaches. It may be important to promote timely dialogue between the sponsors and FDA review divisions to enable proactive management of potential issues, which could become major issues either for submission or review of the premarket application. Additionally, the development of guidance that outlines best practices or novel approaches for avoiding commonly encountered issues may be of value.

Rather than a lengthy briefing document, a proposed Type D meeting (See Table 1 below) may be supported by a brief memo containing the information needed to address the drug development issue(s) at hand to inform the discussion and feedback provided for focused meetings. Additionally, templates that provide high-level summaries of specific aspects of drug development such as the status of CMC development and dose optimization to facilitate early FDA identification of potential drug development issues that could be addressed in future meetings with the sponsor post-BTD might be beneficial (see discussion on use of “Drug Development Snapshots” as a potential communication tool).

During the focus group meetings, there was discussion of informal meeting requests but ultimately the value of a more “informal request” may be limited when there are questions requiring input from multiple review disciplines or limited background is provided by the sponsor. The ability to connect informally with review staff, Division leaders, and Office Directors, while potentially valuable for timely decision making, may be counterproductive unless it includes all relevant members of the FDA review team and is incorporated as official FDA correspondence to the sponsor.

Table 1: Available FDA Meeting Types and Applicability for Products with BTD^{7,8,9}

Meeting Type	General Purpose	Timeframe	Application for BTD Products
Type A Meetings	Meetings that are necessary for an otherwise stalled product development program to proceed or to address an important safety issue.	Within 30 days of request	Strengthening communication between the FDA and sponsor could help avoid issues that would require a Type A meeting for products with BTD.
Type B Meetings	Routine meetings occurring at pre-defined endpoints between FDA and a sponsor. Meetings typically occur right after or right before the submission of clinical data or a new drug filing.	Within 60 to 70 days of request	Formal meetings for products with BTD, including the initial comprehensive BTD meeting, are granted as Type B, unless they qualify as Type A meetings.
Type C Meetings	A meeting that is not a Type A or Type B meeting regarding the development and review of a product.	Within 75 days of request	Very rarely used for BTD products, as BTD default would be to Type B meetings unless reclassified by FDA and have a potentially shorter turnaround. With the addition of a Type D meeting (see below), Type C meetings could be reserved for development issues that would not warrant as quick of a turnaround.
Type F Meetings	Early advice meetings to discuss pediatric development plans.	Within 30 days of request	Meetings encouraged for BTD drugs to ensure that an agreed initial pediatric study plan (iPSP) is in place prior to marketing application. This may be particularly important with respect to mechanism of action based pediatric requirements (Section 504 of FDARA 2017) for original application of oncology drugs.
Proposed Type D Meetings*	Focused on a narrow set of issues (no more than 2 focused topics) which could include: <ul style="list-style-type: none"> a follow-up question that raises a new issue after a formal meeting a narrow issue with few associated questions a general question about an innovative development approach 	Within 50 days of request	This meeting type could promote proactive, collaborative discussion to help identify and mitigate/troubleshoot potential rate-limiting steps to development articulated by the sponsor in a brief memo (as opposed to the typical Q&A longer format seeking agreement from the FDA on specific questions). Type D meetings could be reserved for issues that require more rapid feedback versus issues that may be best routed through a Type B meeting request.

*Meeting type described in the PDUFA VII Commitment Letter (<https://www.fda.gov/media/151712/download>)



Opportunity 3. Develop communication tools and optimize processes to support interdisciplinary coordination within and between the FDA and sponsors.

Create a program for voluntary submission of “Drug Development Snapshots” for earlier identification of issues that could result in delays in development.

A strength of BTM is the product-specific dialogue provided through cross-functional structured interactions between the FDA and sponsors, which involves senior leadership at FDA. However, challenges may be encountered at different times and in different areas of drug development depending on the drug development program. Identification of challenges may not align with milestone meetings that are typically attended by cross-functional groups within the FDA and the drug development team, which can result in delays addressing these issues. Voluntary submission of high-level information in the form of periodic Drug Development Snapshots on certain development topics (see mock template for dosing snapshot in Appendix) or issues by the sponsor could help prompt earlier identification of rate-limiting aspects of development (e.g., dosing, CMC/Product quality aspects, diagnostic co-development, plans for confirmatory trials if accelerated approval pathway is anticipated, etc.), serve as a vehicle to support information exchange, and help determine when meetings outside of the normal milestone cadence would be most beneficial. This increased transparency throughout development outside of meetings can also provide cross-FDA alignment on where the development program stands, plans for upcoming milestones, and necessary interventions. While these snapshots could be particularly useful for enabling improved real-time communication, they could also be leveraged beyond BTM products. Initiating a pilot project to explore the utilization of Drug Development Snapshots, including the timing with respect to drug development of when they would be helpful and understanding the types of information and level of detail included, could be considered.

Refine best practices for communicating with RPMs to help facilitate efficient collaboration.

FDA Regulatory Project Managers (RPMs) play a vital role in triaging and prioritizing sponsors' requests, as well as in identifying the appropriate person(s) for FDA-sponsor and internal meetings, and in coordinating responses for such meetings when necessary. While sponsor interactions with RPMs are extremely helpful, there is opportunity to improve these interactions. Defining the best communication practices for sponsors and RPMs may help sponsors understand expectations specifically in the context of a program with BTM. One opportunity to optimize efficient communication is the ability for sponsors to “flag” requests for feedback that are time-sensitive, and clearly identify they are requesting a reviewer's feedback on a specific topic, to help RPMs appropriately prioritize requests. RPMs could also provide a time estimate for how long it will take to provide feedback for the request, at the point of acknowledgement of the request. Updating CDER's Manual of Policies and Procedures (MAPP) 6030.9 Good Review Practice: Good Review Management Principles and Practices for Effective IND Development and Review, which describes review management principles and practices, may provide additional clarity on best communication practices. A one-on-one meeting between the FDA and Sponsor RPMs could also help set communication expectations. Further, CDER's 2017 Best Practices for Communication Between IND Sponsors and FDA During Drug Development Good Review Practice¹⁰ outlines appropriate communication strategies between sponsors and FDA, and addi-

tional awareness and following of the best practices may increase efficiencies.

Communicate the sponsor's role in preparing materials for cross-discipline meetings.

For cross-discipline meetings in BTM drug development to be productive, it may be helpful to outline the role of both sponsors and the FDA in identifying when general or more specific feedback is warranted and the key disciplines needed for each interaction. Sponsors could provide concise, focused information necessary for FDA to answer the questions at hand. RPMs could then distribute the materials to the review team, including to reviewers or consultants outside of the Division or Center.

Opportunity 4. Provide a roadmap for addressing key pressure points for products with BTM.

Encourage early collaboration, alignment, and prioritization between pharmaceutical and device sponsors and CDER and CDRH.

Challenges to efficient development of a companion diagnostic can arise particularly for drugs developed for rare patient populations or in the setting of a product with BTM. Early identification of the need for a companion diagnostic and plans for parallel development with the goal of contemporaneous approval of a BTM drug and companion diagnostic (if needed) could be a key component of the post-BTM meeting. As noted earlier, outlining specific meeting types (Type B meetings or otherwise) and timelines to focus on incorporating companion diagnostic co-development can increase collaboration between CDER, CDRH, the pharmaceutical sponsor, and the diagnostic sponsor and enable early preparation for possible bridging studies, as well as strategies for saving patient samples and adequate patient ascertainment. Additionally, notification to CDRH upon designation of a BTM could allow for efficient mobilization of appropriate resources. There is also a need for additional clarity around the level of evidence and data elements needed prior to approval for a companion diagnostic. It may be helpful if diagnostic tests developed to direct the use of therapies with BTM were also considered for breakthrough device designation to assist in alignment, prioritization, and collaboration between senior leadership within and across medical product Centers responsible for each breakthrough product.

Facilitate timely discussion and agreement on dose selection, exposure-response analyses, and study design.

Rapid development programs associated with BTM can give the impression that there is insufficient time for robust dose finding approaches; however, identifying the optimal dosage to support safe and effective use of oncology drugs, and accumulating sufficient information to support this dosage, is extremely important, and the selection of the recommended dose without adequate investigation is unacceptable. FDA's OCE has highlighted dose optimization, including for BTM drugs, as a priority by introducing Project Optimus, calling for dose selection justification and earlier discussion of dose selection during the IND phase. BTM may be granted prior to identification of the optimal dose; however, the approach planned to support the dosage(s) intended for further development could be discussed with the FDA prior to embarking on a clinical trial intended to provide evidence of safety and effectiveness to support a marketing application.



These approaches could integrate PK, PD, efficacy, safety, and tolerability data to adequately support dose selection and may result in the selection of 2 or more doses for further exploration. Sponsors could seek out these discussions, which may occur before or after receipt of BTD, as early in the development process as possible and FDA could provide timely dialogue to support dose optimization prior to initiation of the trial intended to support a marketing application. FDA could outline opportunities to discuss strategies for dose optimization and selection of the pivotal dose(s) in pre-BTD meetings. FDA could also clarify and provide feedback on the appropriate use of systems and model-based approaches to support dose selection, study design, and exposure-response analyses and provide feedback on proposals to leverage relevant markers of activity to inform the dosing decisions early in development. Further, learnings from FDA's work through its PDUFA VI commitment on model-informed drug development (MIDD) could be leveraged to facilitate appropriate dose selection. Clarity is needed on the appropriate use of MIDD to support dose selection, including what evidence is needed to show that a model is credible and the role of the model in supporting or supplementing clinical data.

Identify processes to support early Office of Pharmaceutical Quality (OPQ) discussions and facilitate timely submission of CMC information to address rate limiting-steps in the commercialization process.

Sponsors are encouraged to provide available CMC/manufacturing information and commercialization plans, which may be in the form of a "Drug Development Snapshot," early to FDA to potentially maximize the ability to address rate-limiting steps in the development and marketing of a breakthrough product. Currently, discussions with CMC/OPQ generally occur later in development, close to the time of planned submission of a marketing application. It may be helpful to have these conversations earlier in the development program, in a more proactive manner so manufacturing and product quality strategies can be developed likely before data are available to support definitive discussions. An opportunity to engage in a discussion specific to late phase/commercial manufacturing and testing approaches as well as to troubleshoot Quality-related development challenges may expedite the commercialization process. As described in the PDUFA VII Commitment Letter, FDA plans to issue a new MAPP on approaches to address CMC challenges for products with accelerated clinical development timelines and will describe early engagement with sponsors of such products. Further, clarity in the process for identifying the OPQ program manager (OPQPM) is needed. Processes for rolling submission and review of information before all the necessary stability data are available could be explored; for example, this could lead to submission of the entire module 3 except for 3.2.S.7 and 3.2.P.8 sections months earlier, with the two sections submitted in the final wave of the NDA. Further, it may be beneficial to discuss early submission plans with OPQ at the EOP 2 or a Pre-NDA meeting. Early submission of detailed manufacturing site information (e.g., list of manufacturing facilities with addresses and FEI numbers, current CGMP status, facilities' prior experience with similar manufacturing processes, manufacturing area and filling line or equipment used) can allow earlier coordination and planning if a pre-licensure or pre-approval inspection is necessary. Further, clarifying the level of CMC information needed for approval may be helpful as well as the types of evidence that could be provided in the post-market setting.

Consider FDA's available strategies to assess facility risks and enable more efficient inspections.

Certain flexibilities were allowed during the COVID-19 pandemic, including operational processes (both for FDA and sponsors), and clarifications on necessary versus optional application components. Operational flexibilities have included remote evaluations and prioritizing certain inspections over others. Considering whether some of these flexibilities, or the principles behind them, could be explored to ensure that inspections for BTD products are timely thus alleviating delays in approval and enabling sponsors and FDA to allocate resources efficiently.

Explore decoupling drug substance and drug product process performance qualification (PPQ).

When drug substance and drug product PPQ occur sequentially (for those programs requiring PPQ data in the initial application), the PPQ timeline may delay submission of the application and product approval. Inclusion of CMC into Real-Time Oncology Review (RTOR) and the development of the CMC Assessment Aid have brought flexibility and enhanced CMC review efficiency to oncology application reviews. Concurrent execution and completion of the drug substance and drug product PPQs could build on experiences in small molecule development and may result in expedited CMC readiness to meet clinical timelines. Feasibility of this approach has been demonstrated in the development of small molecules over the last few decades as well as for the BTD product pembrolizumab (a monoclonal antibody).¹¹ Exploring how to successfully decouple drug substance PPQ and drug product PPQ, could help to expedite the timeline safely and efficiently. For small molecules, concurrent validation approaches could be useful for a BTD product to market the PPQ batches. Circumstances and rationale for concurrent release could be fully described in a PPQ protocol. More details can be found in the process validation guidance.¹²



References

1. FDA. Guidance for Industry: Expedited Programs for Serious Conditions--Drugs and Biologics. (2014). Available at: <https://www.fda.gov/media/86377/download>. (Accessed: 7th July 2021)
2. FDA. Guidance for Industry: Expedited Programs for Serious Conditions--Drugs and Biologics. (2014). Available at: <https://www.fda.gov/media/86377/download>. (Accessed: 7th July 2021)
3. FDA. CDER Breakthrough Therapy Designation Requests Received by Fiscal Year. (2020). Available at: <https://www.fda.gov/media/95292/download>. (Accessed: 12th August 2021).
4. Friends of Cancer Research. Drug Development Dashboard: Original Approvals in Oncology, 2013-2021. (2021). Available at: <https://friendsofcancerresearch.org/drug-development-dashboard>
5. FDA. Manual of Policies and Procedures for Good Review Practice management of BTB drugs and biologics (2014). Available at: <https://www.fda.gov/files/about%20fda/published/Good-Review-Practice--Management-of-Breakthrough-Therapy-Designated-Drugs-and-Biologics.pdf> (Accessed 29 July 2021).
6. FDA. PDUFA Reauthorization Performance Goals and Procedures Fiscal Years 2023 Through 2027 (2022). Available at: <https://www.fda.gov/media/151712/download> (Accessed 9 September 2021).
7. FDA. Formal Meetings Between the FDA and Sponsors or Applicants of PDUFA Products Guidance for Industry (2017). Available at: <https://www.fda.gov/media/109951/download> (Accessed 29 July 2021).
8. FDA. Manual of Policies and Procedures for Good Review Practice management of BTB drugs and biologics (2014). Available at: <https://www.fda.gov/files/about%20fda/published/Good-Review-Practice--Management-of-Breakthrough-Therapy-Designated-Drugs-and-Biologics.pdf> (Accessed 29 July 2021).
9. FDA. Pediatric Oncology Product Development Early Advice Meeting (Type F)1. Available at: <https://www.fda.gov/about-fda/oncology-center-excellence/pediatric-oncology-product-development-early-advice-meeting-type-f1> (Accessed 29 July 2021).
10. FDA. Best Practices for Communication Between IND Sponsors and FDA During Drug Development Guidance for Industry and Review Staff Good Review Practice (2017). Available at: <https://www.fda.gov/media/94850/download> (Accessed 8 September 2021).
11. Dye, E., Sturgess, A., Maheshwari, G. et al. Examining Manufacturing Readiness for Breakthrough Drug Development. *AAPS PharmSciTech*: 17, 529–538. (2016). <https://doi.org/10.1208/s12249-015-0455-1>
12. FDA. Guidance for Industry Process Validation: General Principles and Practices (2011). Available at: <https://www.fda.gov/files/drugs/published/Process-Validation--General-Principles-and-Practices.pdf> (Accessed 9 September 2021).

Appendix 1. Drug Development Snapshot Template – Clinical Pharmacology (Dose & Administration) Snapshot

Please note: The table below describes the supportive evidence for the proposed dose and schedule. The target length of the completed snapshot is 2–5 pages.

Key Area of Consideration	Supporting Evidence
Recommended dose, schedule and route of administration	<ul style="list-style-type: none"> • What is the current dose(s), schedule(s) and route of administration that are currently being evaluated in clinical trials? Has the RP2D been selected? If the RP2D has not been selected, what key questions are outstanding? • When do you anticipate that a R2PD will be selected? • Are other routes of administration being investigated?
Mechanism of action (MOA) and format	<ul style="list-style-type: none"> • Is the therapeutic a small or large molecule? Another platform? What is the MOA?
Translational evidence	<ul style="list-style-type: none"> • Is there established pharmacological evidence (e.g., target engagement, MOA, outcome-based biomarkers, tumor volume) in the relevant preclinical species? • Is the dose-PK relationship established in the non-clinical species (i.e., is the PK dose proportional)? • Are the pharmacological/efficacious target concentrations for patients defined? • Is the dose/exposure-response (i.e., biomarkers, tumor size, etc.) relationship identified from the in vitro cellular systems or the in vivo animal models?
Clinical Evidence	
Clinical studies	<ul style="list-style-type: none"> • List of ongoing and completed studies (i.e., single agent and/or combination studies, indication, etc.) • Brief description of study design including patient population/cancer type(s) under study, line of therapy, and doses and schedules evaluated, sample size. For example, the following elements can be considered: <ul style="list-style-type: none"> ○ Dose escalation, expansion cohorts with or without randomization ○ Single arm randomization (i.e., dose and/or control); adaptive design

PK characteristics	<ul style="list-style-type: none"> ● Is the dose-PK relationship established (i.e., is the PK dose proportional)? ● Do the PK characteristics (accumulation, half-life) justify the dosing interval? ● Are there any intrinsic or extrinsic factors (e.g., food, body weight, immunogenicity) that would majorly influence PK? (i.e., if these warrants dose adjustments in a subset of patients) ● Was the PK variability considered when selecting a dose that would achieve target exposure for the majority of patients?
Safety summary	<ul style="list-style-type: none"> ● Summary of frequencies of key AEs (including chronic low grade AEs which can affect tolerability) of interest by dose ● Is there a dose/exposure-safety or PK-PD relationship, upon the adjustment of potential covariates, for safety? If yes, what is the nature of the relationship? ● Summary of dose interruptions, reductions, and discontinuations by dose/exposures <ul style="list-style-type: none"> ○ Is there an increased frequency of dose interruptions or reductions or treatment discontinuations with increasing doses/exposures? ● Are there any late occurrence toxicities beyond the DLT period? Are there early PD biomarkers reflective of the delayed safety endpoints? ● Are there any overlapping toxicities with the concomitant medications in the patient population (e.g., treatment combinations for NME with SOC and/or treatments for comorbidities/cancer related symptoms)? ● Is there an increased frequency of dose interruptions, reductions, or treatment discontinuations with increasing doses/exposures? ● If acute/transient toxicities were observed, were alternative dosing approaches considered (e.g., step-up dosing)? ● Does existing data indicate this is a narrow therapeutic window drug with dose limiting toxicity that is monitorable (e.g., biomarkers, BP, HR, neuropathy)? ● If yes, does this drug provide an opportunity to personalize the dose for an individual patient or a sub-population based on the emerging monitorable toxicity?
Efficacy summary	<ul style="list-style-type: none"> ● Summary of response endpoints by dose (e.g., ORR, PFS) ● Is there a dose/exposure – efficacy (primary efficacy endpoint) and PK-PD (e.g., mechanism of action/predictive biomarkers) relationship upon the adjustment of potential confounders? If yes, what is the nature of the relationship? ● Is the dose schedule (e.g., frequency, dose holidays) justified based on the K/PK-PD and/or QSP modeling approaches? ● Are the relevant exposure metrics for efficacy identified (e.g., AUC, C_{max}, C_{min}, concentration-time, RO)?

Other considerations	<ul style="list-style-type: none"> • Are there any manufacturing considerations (e.g., pill burden, maximal feasible dose, etc.) that needs to be taken into account? • Is there any patient factors that need to be considered (e.g., patient convenience/compliance [QD, BID, TID), QW vs Q3W, SC vs IV) • Complimentary M&S approaches (i.e., PK-TGI/QSP/ML etc.) for dose optimization and/or inform dose adjustments
Additional Clinical Evidence	
Planned clinical studies	<ul style="list-style-type: none"> • Are there additional planned clinical studies that will contribute data to the current D&A plan/rationale or future D&A proposals?
Other Evidence	<ul style="list-style-type: none"> • Does additional scientific evidence exist (e.g., from similar class, MOA or indication) that may support the current D&A plan/rationale (e.g., publications, scientific presentations)?
<p>Abbreviations: AEs=adverse events; AUC=area under the curve; BP=blood pressure; C_{max}=maximum 'peak' concentration; C_{min}=minimum 'trough' concentration; D&A=dose & administration; DLT=dose-limiting toxicity; HR=heart rate; K=kinetic; MOA=mechanism of action; NME=new molecular entity; ORR=overall response rate; PD=pharmacodynamic; PFS=progression-free survival; PK=pharmacokinetic; QSP=quantitative systems pharmacology; RO=receptor occupancy; SOC=standard of care</p>	